

PARENT STOCK

ROSS 308 FF

Performance
Objectives

2016

Fast Feathering



Introduction

This booklet contains the performance objectives for Ross® 308 FF parent stock (fast feathering) and should be used in conjunction with the **Ross Parent Stock Management Handbook** and the **Ross 308 FF Management Supplement**.

Performance

Poultry production is a global activity, but across the world there are differing management strategies adapted to local conditions.

These performance objectives are for birds that receive the first light stimulation **after** 21 weeks (147 days) of age. This is the most common strategy used worldwide as it gives distinct advantages in early egg size, chick number, and broiler chick quality. If flocks reach 5-10% production prior to 25 weeks of age early egg size will be reduced, resulting in smaller chicks. In managing this, the timing of photostimulation is key.

Achieving the genetic potential of the birds depends on:

- Management to provide birds with their required environment.
- A dietary regime that provides the appropriate nutrients.
- Effective biosecurity and disease control.

If any one of these elements is sub-optimal, performance will suffer. The 3 sectors, environment, nutrition and health, are also interdependent. A problem in any one will result in a negative response by the bird to the other factors.

Data contained within this booklet indicates the performance that can be achieved under good management and environmental conditions and when feeding the recommended nutrient levels. They should therefore be regarded as “Performance Objectives” and not specifications. In practice, variations in performance may occur for a wide variety of reasons. For example, feed consumption can be affected significantly by form of feed, energy level, and house temperature.

While every attempt has been made to ensure the accuracy and relevance of the information presented, Aviagen® accepts no liability for the consequences of using this information to manage parent stock.

All weight measurements are shown in both **metric (kg/g)** and **imperial (lb/oz)** to reflect the global nature of this publication.

In the tables, values are rounded. This may result in small inaccuracies when using the objectives to calculate other performance statistics.

For more information on the management of Ross stock, please contact your local Ross representative.

Contents

03	Performance Summary
04	Female In-Season Body Weight and Feeding Program
05	Female Out-of-Season Body Weight and Feeding Program
06	Feeding into Lay and Nutrition Allocation at Peak
07	Male Body Weight and Feeding Program
08	Weekly Egg Production
9	Weekly Hatchability and Chick Production
10	Weekly Egg Weight and Egg Mass

Performance Summary

Global Ross 308 FF breeder performance objectives for birds light-stimulated **after** 21 weeks (147 days).

Summary of 40 weeks of production.

Age at depletion (days) (weeks)	448 64	448 64
Total Eggs (HHA*)	185	185
Hatching Eggs (HHA*)	175	175
Chicks/female housed at 175 days (25 weeks)	150	150
Hatchability %	85.8	85.8
Age at 5% Production (days) (weeks)	175 25	175 25
Peak Production %	86.8	86.8
Body weight at 175 days (25 weeks)	2975 g	6.56 lb
Body weight at depletion	4080-4180 g	8.99-9.22 lb
Mortality + culls % (rearing period)	4-5	4-5
Mortality % (laying period)	8	8
Feed/100 Chicks** day old - 448 days (0-64 weeks)	37.0 kg	81.6 lb
Feed/100 Hatching Eggs** day old - 448 days (0-64 weeks)	31.8 kg	70.1 lb

KEY
 (kg/g) – metric measurement
 (lb/oz) – imperial measurement

NOTES

* *Hen-Housed Average.*

** *Feed amounts expressed in the table do not include male feed allocations.*

Female In-Season Body Weight and Feeding Program

All flocks grown in black-out housing are considered in-season.

Age (days)	Age (weeks)	Body Weight (g)	Weekly Gain (g)	Feed (g/bird/day)	Body Weight (lb)	Weekly Gain (lb)	Feed (lb/100/day)	Energy Intake (kcal/bird/day)*
Day old	0	40		ad lib	0.09		ad lib	ad lib
7	1	115	75	24	0.25	0.16	5.2	66
14	2	215	100	28	0.47	0.22	6.2	78
21	3	335	120	32	0.74	0.27	6.9	88
28	4	450	115	35	0.99	0.25	7.7	98
35	5	560	110	38	1.23	0.24	8.4	107
42	6	660	100	41	1.45	0.22	9.0	115
49	7	760	100	45	1.67	0.22	9.8	125
56	8	860	100	48	1.89	0.22	10.5	134
63	9	960	100	50	2.11	0.22	11.0	140
70	10	1060	100	53	2.33	0.22	11.6	148
77	11	1160	100	56	2.56	0.23	12.4	158
84	12	1260	100	60	2.78	0.22	13.2	167
91	13	1360	100	63	3.00	0.22	14.0	177
98	14	1460	100	67	3.22	0.22	14.7	187
105	15	1560	100	71	3.44	0.22	15.6	199
112	16	1670	110	76	3.68	0.24	16.7	213
119	17	1790	120	80	3.94	0.26	17.6	224
126	18	1915	125	86	4.22	0.28	18.9	240
133	19	2050	135	92	4.52	0.30	20.2	257
140	20	2195	145	98	4.83	0.31	21.6	274
147	21	2345	150	105	5.17	0.34	23.1	294
154	22	2500	155	111	5.51	0.34	24.4	310
161	23	2660	160	116	5.86	0.35	25.5	325
168	24	2820	160	122	6.21	0.35	26.8	341
175	25	2975	155	128	6.55	0.34	28.3	359
182	26	3120	145	138	6.87	0.32	30.3	385
189	27	3245	125	151	7.15	0.28	33.3	423
196	28	3340	95	164	7.36	0.21	36.1	459
203	29	3395	55	164	7.48	0.12	36.1	459
210	30	3435	40	164	7.57	0.09	36.1	459
217	31	3465	30	164	7.63	0.06	36.1	459
224	32	3490	25	164	7.69	0.06	36.1	459
231	33	3510	20	164	7.73	0.04	36.1	459
238	34	3530	20	164	7.78	0.05	36.1	459
245	35	3550	20	164	7.82	0.04	36.1	459
252	36	3570	20	164	7.86	0.04	36.1	458
259	37	3590	20	163	7.91	0.05	36.0	457
266	38	3610	20	163	7.95	0.04	35.9	456
273	39	3630	20	163	8.00	0.05	35.8	456
280	40	3650	20	162	8.04	0.04	35.8	455
287	41	3670	20	162	8.08	0.04	35.7	454
294	42	3690	20	162	8.13	0.05	35.6	453
301	43	3710	20	161	8.17	0.04	35.5	452
308	44	3730	20	161	8.22	0.05	35.5	451
315	45	3750	20	161	8.26	0.04	35.4	450
322	46	3770	20	160	8.30	0.04	35.3	449
329	47	3790	20	160	8.35	0.05	35.2	448
336	48	3810	20	160	8.39	0.04	35.2	447
343	49	3830	20	159	8.44	0.05	35.1	446
350	50	3850	20	159	8.48	0.04	35.0	445
357	51	3870	20	159	8.52	0.04	35.0	444
364	52	3890	20	158	8.57	0.05	34.9	443
371	53	3910	20	158	8.61	0.04	34.8	443
378	54	3930	20	158	8.66	0.05	34.7	442
385	55	3950	20	157	8.70	0.04	34.7	441
392	56	3970	20	157	8.74	0.04	34.6	440
399	57	3990	20	157	8.79	0.05	34.5	439
406	58	4010	20	156	8.83	0.04	34.4	438
413	59	4030	20	156	8.88	0.05	34.4	437
420	60	4050	20	156	8.92	0.04	34.3	436
427	61	4070	20	155	8.96	0.04	34.2	435
434	62	4090	20	155	9.01	0.05	34.2	434
441	63	4110	20	155	9.05	0.04	34.1	433
448	64	4130	20	154	9.10	0.05	34.0	432

KEY
 (kg/g) – metric measurement
 (lb/oz) – imperial measurement

North of Equator: Flocks hatched August-December.

South of the Equator: Flocks hatched February-June.

January and July are transitional months so lighting programs for placements during these 2 months should be based on individual experience and location.

*Feed quantities are given as a guide only, based on recommend dietary energy levels of a 2- or 3-stage rearing program (2800 kcal ME/kg; 1270 kcal ME/lb). Adjustments must be made to reflect feeding differing energy levels.

NOTES
 Weekly body-weight gain beyond 33 weeks (231 days) should average approximately 20 g (0.04-0.05 lb).
 Body weights are based on a feed day, 4-6 hours after feeding.

Female Out-of-Season Body Weight and Feeding Program

Age (days)	Age (weeks)	Body Weight (g)	Weekly Gain (g)	Feed (g/bird/day)	Body Weight (lb)	Weekly Gain (lb)	Feed (lb/100/day)	Energy Intake (kcal/bird/day)*
Day old	0	40		ad lib	0.09		ad lib	ad lib
7	1	115	75	23	0.25	0.16	5.2	66
14	2	215	100	28	0.47	0.22	6.2	79
21	3	330	115	32	0.73	0.26	7.0	89
28	4	450	120	35	0.99	0.26	7.8	99
35	5	560	110	39	1.23	0.24	8.5	109
42	6	660	100	42	1.45	0.22	9.2	117
49	7	760	100	45	1.67	0.22	10.0	127
56	8	870	110	49	1.92	0.25	10.7	136
63	9	980	110	51	2.16	0.24	11.2	143
70	10	1090	110	54	2.40	0.24	11.9	151
77	11	1200	110	58	2.64	0.24	12.8	162
84	12	1300	100	62	2.86	0.22	13.6	172
91	13	1400	100	66	3.08	0.22	14.5	184
98	14	1500	100	70	3.30	0.22	15.4	195
105	15	1610	110	75	3.55	0.25	16.4	209
112	16	1740	130	80	3.83	0.28	17.7	225
119	17	1880	140	85	4.14	0.31	18.6	237
126	18	2020	140	90	4.45	0.31	19.9	253
133	19	2160	140	96	4.76	0.31	21.0	268
140	20	2300	140	101	5.07	0.31	22.2	282
147	21	2460	160	106	5.42	0.35	23.4	298
154	22	2640	180	111	5.81	0.39	24.5	312
161	23	2800	160	116	6.17	0.36	25.5	325
168	24	2950	150	122	6.50	0.33	26.8	341
175	25	3090	140	129	6.81	0.31	28.3	360
182	26	3220	130	138	7.09	0.28	30.5	388
189	27	3330	110	152	7.33	0.24	33.6	426
196	28	3420	90	166	7.53	0.20	36.5	464
203	29	3490	70	166	7.69	0.16	36.5	464
210	30	3540	50	166	7.80	0.11	36.5	464
217	31	3580	40	166	7.89	0.09	36.5	464
224	32	3610	30	166	7.95	0.06	36.5	464
231	33	3630	20	166	8.00	0.05	36.5	464
238	34	3650	20	166	8.04	0.04	36.5	464
245	35	3670	20	166	8.08	0.04	36.5	464
252	36	3690	20	165	8.13	0.05	36.4	463
259	37	3710	20	165	8.17	0.04	36.4	462
266	38	3730	20	165	8.22	0.05	36.3	461
273	39	3750	20	164	8.26	0.04	36.2	460
280	40	3770	20	164	8.30	0.04	36.1	459
287	41	3790	20	164	8.35	0.05	36.1	458
294	42	3810	20	163	8.39	0.04	36.0	458
301	43	3830	20	163	8.44	0.05	35.9	457
308	44	3850	20	163	8.48	0.04	35.8	456
315	45	3870	20	162	8.52	0.04	35.8	455
322	46	3890	20	162	8.57	0.05	35.7	454
329	47	3910	20	162	8.61	0.04	35.6	453
336	48	3930	20	161	8.66	0.05	35.6	452
343	49	3950	20	161	8.70	0.04	35.5	451
350	50	3970	20	161	8.74	0.04	35.4	450
357	51	3990	20	160	8.79	0.05	35.3	449
364	52	4010	20	160	8.83	0.04	35.3	448
371	53	4030	20	160	8.88	0.05	35.2	447
378	54	4050	20	159	8.92	0.04	35.1	446
385	55	4070	20	159	8.96	0.04	35.0	445
392	56	4090	20	159	9.01	0.05	35.0	444
399	57	4110	20	158	9.05	0.04	34.9	443
406	58	4130	20	158	9.10	0.05	34.8	443
413	59	4150	20	158	9.14	0.04	34.7	442
420	60	4170	20	157	9.19	0.05	34.7	441
427	61	4190	20	157	9.23	0.04	34.6	440
434	62	4210	20	157	9.27	0.04	34.5	439
441	63	4230	20	156	9.32	0.05	34.4	438
448	64	4250	20	156	9.36	0.04	34.4	437

KEY

- (kg/g) – metric measurement
- (lb/oz) – imperial measurement

North of Equator: Flocks hatched February - June.

South of the Equator: Flocks hatched August - December.

January and July are transitional months so lighting programs for placements during these 2 months should be based on individual experience and location.

*Feed quantities are given as a guide only, based on recommend dietary energy levels of a 2- or 3-stage rearing program (2800 kcal ME/kg; 1270 kcal ME/lb). Adjustments must be made to reflect feeding differing energy levels.

NOTES

Weekly body-weight gain beyond 33 weeks (231 days) should average approximately 20 g (0.04-0.05 lb).

Body weights are based on a feed day, 4-6 hours after feeding.

Female In-Season Feeding into Lay

Hen-Day (%)	Daily Energy Intake (kcal ME/bird/day)*	Feed Intake (g/bird/day)	Feed Increase (g/bird/day)
5	359	128	
10	367	131	3
15	374	134	3
20	382	136	2
25	387	138	2
30	393	141	3
35	400	143	2
40	406	145	2
45	412	147	2
50	419	149	2
55	425	152	3
60	434	155	3
65	442	158	3
70	450	161	3
peak	459	164	3

Female Out-of-Season Feeding into Lay

Hen-Day (%)	Daily Energy Intake (kcal ME/bird/day)*	Feed Intake (g/bird/day)	Feed Increase (g/bird/day)
5	360	129	
10	367	131	2
15	373	133	2
20	380	136	3
25	390	139	3
30	396	142	3
35	403	144	2
40	409	146	2
45	415	148	2
50	421	151	3
55	428	153	2
60	437	156	3
65	446	159	3
70	455	162	3
peak	464	166	4

*Daily energy and feed intakes are based on current recommended dietary levels of energy (2800 kcal ME/kg; 1270 kcal ME/lb) and assuming an ambient temperature of 20-21°C (68-70°F).

NOTES

Feeding programs should be adjusted according to actual feed intake at 5% hen-day production. It may be necessary to adjust feed amounts daily (rather than every 5% as given in the table), taking into account the rate of daily production. Adjustments to feed amounts will need to be made if dietary energy levels are different to those recommended or if environmental temperatures are warmer or cooler than assumed here.

Female In-Season Nutrient Allocation at Peak

Nutrient	Nutrient Allocation at Peak
Energy (kcal/bird/day)*	459
Digestible Amino Acids (mg/bird/day)	
Lysine	984
Methionine & Cystine	968
Methionine	607
Threonine	804
Valine	918
Isoleucine	820
Argenine	1296
Tryptophan	230
Minerals (mg/bird/day)	
Calcium	4920
Available Phosphorus	574

Female Out-of-Season Nutrient Allocation at Peak

Nutrient	Nutrient Allocation at Peak
Energy (kcal/bird/day)*	464
Digestible Amino Acids (mg/bird/day)	
Lysine	996
Methionine & Cystine	979
Methionine	614
Threonine	813
Valine	930
Isoleucine	830
Argenine	1311
Tryptophan	232
Minerals (mg/bird/day)	
Calcium	4950
Available Phosphorus	581

*Based on a recommended energy level of 2800 kcal ME/kg (1270 kcal ME/lb).

Male Body Weight and Feeding Program

Age (days)	Age (weeks)	Body Weight (g)	Weekly Gain (g)	Feed (g/bird/day)	Body Weight (lb)	Weekly Gain (lb)	Feed (lb/100/day)	Energy Intake (kcal/bird/day)*
Day old	0	40		ad lib	0.09		ad lib	
7	1	150	110	35	0.33	0.24	7.6	97
14	2	320	170	42	0.70	0.37	9.3	118
21	3	525	205	48	1.16	0.46	10.5	134
28	4	755	230	52	1.66	0.50	11.5	147
35	5	945	190	56	2.08	0.42	12.4	158
42	6	1130	185	60	2.49	0.41	13.2	168
49	7	1280	150	63	2.82	0.33	13.9	177
56	8	1420	140	66	3.13	0.31	14.6	185
63	9	1545	125	69	3.40	0.27	15.2	194
70	10	1670	125	72	3.68	0.28	15.9	202
77	11	1795	125	75	3.95	0.27	16.5	210
84	12	1920	125	78	4.23	0.28	17.2	218
91	13	2045	125	81	4.50	0.27	17.8	227
98	14	2170	125	84	4.78	0.28	18.6	236
105	15	2295	125	88	5.06	0.28	19.3	246
112	16	2420	125	92	5.33	0.27	20.2	257
119	17	2560	140	96	5.64	0.31	21.2	269
126	18	2715	155	101	5.98	0.34	22.2	282
133	19	2875	160	106	6.33	0.35	23.3	296
140	20	3035	160	111	6.69	0.36	24.4	310
147	21	3195	160	115	7.04	0.35	25.4	323
154	22	3355	160	120	7.39	0.35	26.3	335
161	23	3515	160	123	7.74	0.35	27.2	346
168	24	3675	160	127	8.09	0.35	27.9	355
175	25	3825	150	134	8.43	0.34	28.4	361
182	26	3960	135	136	8.72	0.29	29.9	366
189	27	4035	75	137	8.89	0.17	30.2	371
196	28	4090	55	139	9.01	0.12	30.5	374
203	29	4120	30	140	9.07	0.06	30.8	377
210	30	4150	30	141	9.14	0.07	31.0	380
217	31	4180	30	141	9.21	0.07	31.2	382
224	32	4210	30	142	9.27	0.06	31.3	384
231	33	4240	30	143	9.34	0.07	31.5	386
238	34	4270	30	144	9.41	0.07	31.6	388
245	35	4300	30	144	9.47	0.06	31.8	389
252	36	4330	30	145	9.54	0.07	31.9	391
259	37	4360	30	145	9.60	0.06	32.0	392
266	38	4390	30	146	9.67	0.07	32.1	394
273	39	4420	30	146	9.74	0.07	32.2	395
280	40	4450	30	147	9.80	0.06	32.3	397
287	41	4480	30	147	9.87	0.07	32.5	398
294	42	4510	30	148	9.93	0.06	32.6	399
301	43	4540	30	148	10.00	0.07	32.7	401
308	44	4570	30	149	10.07	0.07	32.8	402
315	45	4600	30	149	10.13	0.06	32.9	403
322	46	4630	30	150	10.20	0.07	33.0	404
329	47	4660	30	150	10.26	0.06	33.1	406
336	48	4690	30	151	10.33	0.07	33.2	407
343	49	4720	30	151	10.40	0.07	33.3	408
350	50	4750	30	152	10.46	0.06	33.4	410
357	51	4780	30	152	10.53	0.07	33.5	411
364	52	4810	30	153	10.59	0.06	33.6	412
371	53	4840	30	153	10.66	0.07	33.7	413
378	54	4870	30	154	10.73	0.07	33.8	415
385	55	4900	30	154	10.79	0.06	33.9	416
392	56	4930	30	155	10.86	0.07	34.0	417
399	57	4960	30	155	10.93	0.07	34.1	419
406	58	4990	30	155	10.99	0.06	34.2	420
413	59	5020	30	156	11.06	0.07	34.3	421
420	60	5050	30	156	11.12	0.06	34.5	422
427	61	5080	30	157	11.19	0.07	34.6	424
434	62	5110	30	157	11.26	0.07	34.7	425
441	63	5140	30	158	11.32	0.06	34.8	426
448	64	5170	30	158	11.39	0.07	34.9	427

KEY
 (kg/g) – metric measurement
 (lb/oz) – imperial measurement

*Feed quantities are given as a guide only, based on recommended dietary energy levels of a 2- or 3-stage rearing program (2800 kcal ME/kg; 1270 kcal ME/lb) and a male diet in lay (2700 kcal ME/kg; 1225 kcal ME/lb). Adjustments must be made to reflect feeding differing energy levels.

NOTES
 Body weights are those 4-6 hours after feeding.
 This profile allows the male to reach sexual maturity by first egg. Weekly body-weight gain beyond 29 weeks (203 days) should average approximately 30 g (0.06-0.07 lb).
 Field performance has shown that this practice ensures that the body condition of the males is not compromised so they will maintain the best possible fertility levels.

Weekly Egg Production

Week of Production	Age (days)	Age (weeks)	Hen-Housed (%)	Hen-Week (%)*	Eggs/Bird/Week Hen-Housed	Eggs/Bird/Cum. Hen-Housed	Hatching Eggs/Bird/Week**	Hatching Eggs/Bird/Cum.	Hatching Egg Utilization Weekly	Hatching Egg Utilization Cum.
1	175	25	6.5	6.5	0.46	0.46				
2	182	26	23.2	23.3	1.63	2.08	1.12	1.12	68.84	53.75
3	189	27	53.2	53.6	3.73	5.81	3.22	4.34	86.40	74.69
4	196	28	74.7	75.3	5.23	11.04	4.72	9.06	90.30	82.08
5	203	29	83.2	84.1	5.83	16.86	5.42	14.48	93.02	85.86
6	210	30	86.1	87.1	6.03	22.89	5.72	20.20	94.91	88.24
7	217	31	86.8	88.0	6.08	28.97	5.82	26.02	95.77	89.82
8	224	32	86.1	87.5	6.03	35.00	5.82	31.84	96.57	90.98
9	231	33	85.0	86.5	5.95	40.94	5.74	37.58	96.52	91.79
10	238	34	83.8	85.5	5.87	46.81	5.65	43.23	96.30	92.35
11	245	35	82.7	84.5	5.79	52.60	5.57	48.80	96.25	92.78
12	252	36	81.5	83.5	5.71	58.30	5.49	54.29	96.20	93.12
13	259	37	80.4	82.5	5.63	63.93	5.41	59.70	96.14	93.38
14	266	38	79.2	81.5	5.55	69.48	5.32	65.02	95.91	93.59
15	273	39	78.1	80.5	5.47	74.94	5.24	70.26	95.85	93.75
16	280	40	76.8	79.4	5.38	80.32	5.15	75.41	95.78	93.89
17	287	41	75.7	78.3	5.30	85.62	5.07	80.48	95.71	94.00
18	294	42	74.5	77.3	5.22	90.83	4.99	85.47	95.64	94.09
19	301	43	73.4	76.3	5.14	95.97	4.91	90.38	95.57	94.17
20	308	44	72.2	75.3	5.06	101.03	4.83	95.21	95.50	94.24
21	315	45	71.1	74.2	4.98	106.01	4.75	99.96	95.43	94.30
22	322	46	70.0	73.2	4.90	110.90	4.67	104.63	95.36	94.34
23	329	47	68.8	72.1	4.82	115.72	4.59	109.22	95.29	94.38
24	336	48	67.5	70.9	4.73	120.45	4.50	113.72	95.21	94.41
25	343	49	66.4	69.9	4.65	125.09	4.42	118.14	95.14	94.44
26	350	50	65.2	68.8	4.57	129.66	4.34	122.48	95.06	94.46
27	357	51	64.1	67.8	4.49	134.15	4.26	126.74	94.99	94.48
28	364	52	63.0	66.7	4.41	138.55	4.18	130.93	94.91	94.49
29	371	53	61.8	65.6	4.33	142.88	4.10	135.03	94.83	94.50
30	378	54	60.7	64.5	4.25	147.13	4.02	139.05	94.75	94.51
31	385	55	59.5	63.5	4.17	151.29	3.94	143.00	94.67	94.52
32	392	56	58.2	62.2	4.08	155.37	3.86	146.85	94.59	94.52
33	399	57	57.1	61.1	4.00	159.37	3.78	150.63	94.50	94.52
34	406	58	56.0	60.0	3.92	163.29	3.70	154.33	94.42	94.52
35	413	59	54.8	58.9	3.84	167.12	3.62	157.95	94.33	94.51
36	420	60	53.7	57.8	3.76	170.88	3.54	161.49	94.24	94.51
37	427	61	52.5	56.7	3.68	174.56	3.46	164.95	94.16	94.50
38	434	62	51.4	55.6	3.60	178.15	3.38	168.34	94.06	94.49
39	441	63	50.2	54.5	3.52	181.67	3.30	171.64	93.97	94.48
40	448	64	49.0	53.2	3.43	185.10	3.22	174.86	93.96	94.47

NOTES

* Hen-week (%) is based on the assumption that mortality in lay is 8% with 0.2% mortality per week.

** A hatching egg is considered to be an egg which is 50 g (21.2 oz/dozen) or heavier.

Weekly Hatchability and Chick Production

Week of Production	Age (days)	Age (weeks)	Hatch All Eggs (%)*	Cum. Hatchability (%)	Chicks/Week Hen-Housed	Cum. Chicks Hen-Housed
1	175	25				
2	182	26	78.8	78.8	0.88	0.88
3	189	27	81.6	80.8	2.63	3.51
4	196	28	84.0	82.5	3.96	7.47
5	203	29	86.0	83.8	4.66	12.13
6	210	30	87.7	84.9	5.01	17.15
7	217	31	89.0	85.8	5.18	22.33
8	224	32	90.1	86.6	5.24	27.57
9	231	33	90.9	87.3	5.22	32.79
10	238	34	91.5	87.8	5.17	37.96
11	245	35	91.9	88.3	5.12	43.09
12	252	36	92.1	88.7	5.06	48.14
13	259	37	92.2	89.0	4.99	53.13
14	266	38	92.1	89.3	4.90	58.03
15	273	39	91.9	89.4	4.82	62.85
16	280	40	91.6	89.6	4.72	67.56
17	287	41	91.2	89.7	4.62	72.18
18	294	42	90.7	89.8	4.53	76.71
19	301	43	90.2	89.8	4.43	81.14
20	308	44	89.6	89.8	4.33	85.47
21	315	45	89.0	89.7	4.23	89.70
22	322	46	88.4	89.7	4.13	93.83
23	329	47	87.8	89.6	4.03	97.85
24	336	48	86.8	89.5	3.91	101.76
25	343	49	85.8	89.3	3.79	105.55
26	350	50	84.8	89.2	3.68	109.23
27	357	51	83.8	89.0	3.57	112.81
28	364	52	82.8	88.8	3.46	116.27
29	371	53	81.9	88.6	3.36	119.63
30	378	54	80.9	88.4	3.25	122.89
31	385	55	79.9	88.1	3.15	126.04
32	392	56	78.9	87.9	3.04	129.08
33	399	57	77.9	87.6	2.94	132.02
34	406	58	76.9	87.4	2.84	134.87
35	413	59	75.9	87.1	2.75	137.62
36	420	60	75.0	86.9	2.65	140.27
37	427	61	74.0	86.6	2.56	142.83
38	434	62	73.0	86.3	2.47	145.30
39	441	63	72.0	86.0	2.38	147.68
40	448	64	71.0	85.8	2.29	149.97

NOTES

* Hatchability is based on an average egg age of 3 days. Hatchability will drop by 0.5% per day of storage between 7 and 11 days.

Weekly Egg Weight and Egg Mass

Week of Production	Age (days)	Age (weeks)	Hen-Week (%)	Egg Weight (g)	Egg Mass* (g)	Egg Weight (oz/dozen)
1	175	25	6.5	49.4	3.2	20.9
2	182	26	23.3	51.5	12.0	21.8
3	189	27	53.6	52.9	28.3	22.4
4	196	28	75.3	54.1	40.7	22.9
5	203	29	84.1	55.2	46.4	23.4
6	210	30	87.1	56.2	49.0	23.8
7	217	31	88.0	57.0	50.2	24.1
8	224	32	87.5	57.6	50.4	24.4
9	231	33	86.5	58.1	50.3	24.6
10	238	34	85.5	58.6	50.1	24.8
11	245	35	84.5	59.1	50.0	25.0
12	252	36	83.5	59.6	49.8	25.2
13	259	37	82.5	60.0	49.5	25.4
14	266	38	81.5	60.4	49.2	25.6
15	273	39	80.5	60.9	49.0	25.8
16	280	40	79.4	61.3	48.7	26.0
17	287	41	78.3	61.7	48.3	26.1
18	294	42	77.3	62.1	48.0	26.3
19	301	43	76.3	62.4	47.6	26.4
20	308	44	75.3	62.8	47.3	26.6
21	315	45	74.2	63.2	46.9	26.8
22	322	46	73.2	63.6	46.6	26.9
23	329	47	72.1	63.9	46.1	27.1
24	336	48	70.9	64.3	45.6	27.2
25	343	49	69.9	64.5	45.1	27.3
26	350	50	68.8	65.0	44.8	27.5
27	357	51	67.8	65.2	44.2	27.6
28	364	52	66.7	65.6	43.8	27.8
29	371	53	65.6	65.8	43.2	27.9
30	378	54	64.5	66.1	42.7	28.0
31	385	55	63.5	66.4	42.2	28.1
32	392	56	62.2	66.7	41.5	28.2
33	399	57	61.1	67.0	41.0	28.4
34	406	58	60.0	67.2	40.4	28.5
35	413	59	58.9	67.6	39.9	28.6
36	420	60	57.8	67.8	39.2	28.7
37	427	61	56.7	67.9	38.5	28.7
38	434	62	55.6	68.1	37.9	28.8
39	441	63	54.5	68.2	37.2	28.9
40	448	64	53.2	68.3	36.4	28.9

KEY

- (kg/g) – metric measurement
- (lb/oz) – imperial measurement

NOTE

* Egg mass (g) = $\frac{\text{Hen-week (\%)} \times \text{Egg weight (g)}}{100}$

100



www.aviagen.com

Aviagen and the Aviagen logo, and Ross and the Ross logo are registered trademarks of Aviagen in the US and other countries. All other trademarks or brands are registered by their respective owners.

© 2016 Aviagen.

0616-AVNR-063

Fast Feathering