



An Aviagen Brand

Arbor Acres PLUS

Parent Stock Nutrition Specifications

2021

Fast Feathering



- 2 Introduction
- 3 4-Stage Rearing Program
- 4 5-Stage Rearing Program
- 5 Female Nutrient Allocation at Peak Production
- 6 Male Program

Introduction

This booklet contains the nutritional recommendations for Arbor Acres® Plus (fast feathering) parent stock and is to be used with the **Arbor Acres Parent Stock Management Handbook**, the **Arbor Acres Plus Management Supplement**, and the **Arbor Acres Plus Parent Stock Performance Objectives**.

PERFORMANCE

To achieve optimal reproductive performance, it is important that the body-weight profiles recommended in the **Arbor Acres Plus Parent Stock Performance Objectives** are followed. For the nutritional recommendations that follow, nutrient specifications presented have been based upon daily energy allocations that enable body-weight profiles and reproductive performance objectives to be achieved.

Recommendations included in this booklet suggest different rearing programs for the following scenarios:

- **4-Stage Rearing Program** - where a smooth energy transition is applied between rearing and laying phases.
- **5-Stage Rearing Program** - where a developer ration is introduced to smooth the transition to a pre-breeder.
- **Separate Male Feed** - only for males in production.

Please note, these nutrient specifications are based on a common dietary energy level of 2800 kcal/kg (1271 kcal/lb), which must be adapted according to local environmental conditions, ingredient quality and availability, and feeding strategies. Thus, nutrient values must be adjusted proportionally to reflect the feeding of different energy levels, which is especially important when considering digestible lysine. Feed allocation should be determined by body weight, evaluation of fleshing and egg production, and therefore altered to maintain the recommended weight and egg production profiles.

Feed allocations provided in the **Arbor Acres Plus Parent Stock Performance Objectives** should be adjusted proportionally to any change in the energy density. Feed volume is an important tool that can be used to lengthen feed clean-up times and prevent body-weight uniformity loss in the rearing period even when multiple grading sessions are adopted. Feeding a lower dietary energy density Pullet Grower can be achieved using a combination of diluent ingredients (some examples include wheat bran or middlings, rice mill-feed, rice, oat or soy hulls, and inert mineral clay sources such as aluminum silicates). It is crucial to closely monitor feed clean-up times to ascertain that all pullets receive their fair share of feed to maintain good body-weight uniformity.

The energy values used in these specifications are based on assays for Metabolizable Energy (ME) published by the World's Poultry Science Association (WPSA). The values for amino acid digestibility are based on Standardized Ileal Digestibility (SID) assays.

It may be beneficial to use a specific diet for males during the production period. A specification for a male diet is provided in this booklet.

www.aviagen.com

Arbor Acres Plus Parent Stock Nutrition Specifications

Female Parent Stock Nutrient Specifications 4-Stage Rearing Program

		STARTER 1	STARTER 2	GROWER	PRE-BREEDER	BREEDER 1	BREEDER 2	BREEDER 3	
Age fed	days	0-21 days	22-42 days	43-105 days	106 days to 5% production	>5% production to 224 days	225-350 days	After 351 days	
Energy per kg*	kcal	2800	2800	2800	2800	2800	2800	2800	
	MJ	11.7	11.7	11.7	11.7	11.7	11.7	11.7	
Energy per lb	kcal	1271	1271	1271	1271	1271	1271	1271	
DIGESTIBLE AMINO ACIDS									
Lysine (max)**	%	1.00	0.72	0.52	0.49	0.62	0.56	0.52	
Methionine	%	0.46	0.37	0.36	0.34	0.38	0.35	0.34	
Methionine & Cystine	%	0.84	0.68	0.62	0.59	0.62	0.57	0.55	
Threonine	%	0.70	0.60	0.52	0.50	0.55	0.53	0.51	
Valine	%	0.81	0.72	0.60	0.57	0.64	0.60	0.56	
Tryptophan	%	0.18	0.18	0.15	0.15	0.15	0.14	0.13	
Arginine	%	1.15	0.92	0.78	0.75	0.85	0.82	0.79	
Leucine	%	1.20	1.03	0.82	0.79	0.95	0.90	0.86	
Isoleucine	%	0.70	0.58	0.47	0.44	0.52	0.50	0.49	
Histidine	%	0.43	0.32	0.26	0.22	0.30	0.28	0.26	
Crude Protein (min)	%	19.0	17.0	14.0	14.0	15.0	14.0	13.0	
MINERALS									
Calcium	%	1.05	0.94	0.90	1.20	3.00	3.20	3.40	
Available Phosphorus	%	0.50	0.47	0.45	0.45	0.36	0.34	0.32	
Sodium	%	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	
Chloride	%	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	
Potassium	%	0.60-0.90	0.60-0.90	0.60-0.90	0.60-0.90	0.70-0.90	0.65-0.90	0.60-0.90	
ADDED TRACE MINERALS PER KG									
Copper	mg	16				16			
Iodine	mg	2				3			
Iron	mg	40				50			
Manganese	mg	120				120			
Selenium	mg	0.3				0.3			
Zinc	mg	120				120			
ADDED VITAMINS PER KG									
Vitamin A	IU	13000				15000			
Vitamin D3	IU	4000				5000			
Vitamin E	IU	100				130			
Vitamin K (Menadione)	mg	6				9			
Thiamin (B1)	mg	5				6			
Riboflavin (B2)	mg	15				20			
Niacin	mg	50				70			
Pantothenic Acid	mg	20				25			
Pyridoxine (B6)	mg	5				8			
Biotin	mg	0.3				0.6			
Folic Acid	mg	3				5			
Vitamin B12	mg	0.05				0.07			
MINIMUM SPECIFICATION									
Choline per kg	mg	1400				1600			
Linoleic Acid	%	1.25				2.00			

* Energy base value. Nutrients should be factored accordingly when feeding different energy values.

** In order to achieve the amino acid requirements without exceeding the recommended levels of digestible lysine it may be necessary to adopt more complex diets.

Notes: These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.

Arbor Acres Plus Parent Stock Nutrition Specifications

Female Parent Stock Nutrient Specifications 5-Stage Rearing Program

		STARTER 1	STARTER 2	GROWER	DEVELOPER	PRE-BREEDER	BREEDER 1	BREEDER 2	BREEDER 3
Age fed	days	0-21 days	22-42 days	43-105 days	106-140 days	141 days to 5% production	>5% production to 224 days	225-350 days	After 351 days
Energy per kg*	kcal	2800	2800	2800	2800	2800	2800	2800	2800
	MJ	11.7	11.7	11.7	11.7	11.7	11.7	11.7	11.7
Energy per lb	kcal	1271	1271	1271	1271	1271	1271	1271	1271
DIGESTIBLE AMINO ACIDS									
Lysine (max)**	%	1.00	0.72	0.52	0.50	0.48	0.62	0.56	0.52
Methionine	%	0.46	0.37	0.36	0.34	0.34	0.38	0.35	0.34
Methionine & Cystine	%	0.84	0.68	0.62	0.60	0.58	0.62	0.57	0.55
Threonine	%	0.70	0.60	0.52	0.50	0.49	0.55	0.53	0.51
Valine	%	0.81	0.72	0.60	0.58	0.56	0.64	0.60	0.56
Tryptophan	%	0.18	0.18	0.15	0.15	0.15	0.15	0.14	0.13
Arginine	%	1.15	0.92	0.78	0.76	0.74	0.85	0.82	0.79
Leucine	%	1.20	1.03	0.82	0.80	0.78	0.95	0.90	0.86
Isoleucine	%	0.70	0.58	0.47	0.45	0.43	0.52	0.50	0.49
Histidine	%	0.43	0.32	0.26	0.23	0.20	0.30	0.28	0.26
Crude Protein (min)	%	19.0	17.0	14.0	14.0	14.0	15.0	14.0	13.0
MINERALS									
Calcium	%	1.05	0.94	0.90	0.90	1.50	3.00	3.20	3.40
Available Phosphorus	%	0.50	0.47	0.45	0.45	0.35	0.36	0.34	0.32
Sodium	%	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23
Chloride	%	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23	0.18-0.23
Potassium	%	0.60-0.90	0.60-0.90	0.60-0.90	0.60-0.90	0.60-0.90	0.70-0.90	0.65-0.90	0.60-0.90
ADDED TRACE MINERALS PER KG									
Copper	mg			16				16	
Iodine	mg			2				3	
Iron	mg			40				50	
Manganese	mg			120				120	
Selenium	mg			0.3				0.3	
Zinc	mg			120				120	
ADDED VITAMINS PER KG									
Vitamin A	IU			13000				15000	
Vitamin D3	IU			4000				5000	
Vitamin E	IU			100				130	
Vitamin K (Menadione)	mg			6				9	
Thiamin (B1)	mg			5				6	
Riboflavin (B2)	mg			15				20	
Niacin	mg			50				70	
Pantothenic Acid	mg			20				25	
Pyridoxine (B6)	mg			5				8	
Biotin	mg			0.3				0.6	
Folic Acid	mg			3				5	
Vitamin B12	mg			0.05				0.07	
MINIMUM SPECIFICATION									
Choline per kg	mg			1400				1600	
Linoleic Acid	%			1.25				2.00	

* Energy base value. Nutrients should be factored accordingly when feeding different energy values.

** In order to achieve the amino acid requirements without exceeding the recommended levels of digestible lysine it may be necessary to adopt more complex diets.

Notes: These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.

Female Parent Stock Nutrient Specifications Nutrient Allocation at Peak Production

Female In-Season

Nutrient	Nutrient Allocation at Peak
Energy (kcal/bird/day)	467
Digestible Amino Acids (mg/bird/day)	
Lysine	1034
Methionine	634
Methionine & Cystine	1034
Threonine	917
Valine	1067
Tryptophan	250
Arginine	1418
Leucine	1584
Isoleucine	867
Histidine	500
Minerals (mg/bird/day)	
Calcium	5004
Available Phosphorus	600

Female Out-of-Season

Nutrient	Nutrient Allocation at Peak
Energy (kcal/bird/day)	473
Digestible Amino Acids (mg/bird/day)	
Lysine	1047
Methionine	642
Methionine & Cystine	1047
Threonine	929
Valine	1081
Tryptophan	253
Arginine	1436
Leucine	1605
Isoleucine	878
Histidine	507
Minerals (mg/bird/day)	
Calcium	5068
Available Phosphorus	608

Male Parent Stock Nutrient Specifications Separate Diet in Production

		MALE FEED
Age Fed		after 175 days
Energy per kg*	kcal	2800
	MJ	11.7
Energy per lb	kcal	1271
DIGESTIBLE AMINO ACIDS		
Lysine**	%	0.35
Methionine	%	0.33
Methionine & Cystine	%	0.58
Threonine	%	0.43
Valine	%	0.47
Tryptophan	%	0.15
Arginine	%	0.68
Leucine	%	0.66
Isoleucine	%	0.41
Histidine	%	0.16
Crude Protein	%	12.0
MINERALS		
Calcium	%	0.70
Available Phosphorus	%	0.35
Sodium	%	0.18-0.20
Chloride	%	0.20-0.23
Potassium	%	0.60-0.75
ADDED TRACE MINERALS PER KG		
Copper	mg	16
Iodine	mg	2
Iron	mg	40
Manganese	mg	120
Selenium	mg	0.3
Zinc	mg	120
ADDED VITAMINS PER KG		
Vitamin A	IU	13000
Vitamin D3	IU	4000
Vitamin E	IU	100
Vitamin K (Menadione)	mg	6
Thiamin (B1)	mg	5
Riboflavin (B2)	mg	15
Niacin	mg	50
Pantothenic Acid	mg	20
Pyridoxine (B6)	mg	5
Biotin	mg	0.3
Folic Acid	mg	3
Vitamin B12	mg	0.05
MINIMUM SPECIFICATION		
Choline per kg	mg	1400
Linoleic Acid	%	1.25

* Energy base value. Nutrients should be factored accordingly when feeding different energy values.

** In order to achieve the amino acid requirements without exceeding the recommended levels of digestible lysine it may be necessary to adopt more complex diets.

Notes: These feed specifications should be used as a guide. They may require adjustment for local conditions, legislation and markets.



Aviagen and the Aviagen logo, and Arbor Acres and the Arbor Acres logo are registered trademarks of Aviagen in the US and other countries. All other trademarks or brands are registered by their respective owners.

Privacy Policy: Aviagen collects data to effectively communicate and provide information to you about our products and our business. This data may include your email address, name, business address and telephone number.

To view the full Aviagen privacy policy visit [Aviagen.com](http://www.aviagen.com).

© 2021 Aviagen.

0421-AVNAA-049